

Handy Handouts®

Free informational handouts for educators, parents, and students

Understanding Childhood Apraxia

By Becky Spivey, M.Ed.



Childhood Apraxia of Speech (CAS) is a very specific motor speech disorder. Children with this diagnosis have difficulty planning and consistently producing sequences of speech movements using their tongue, lips, jaw, and palate. These difficulties limit a listener's understanding of the child's speech and prevent the child from producing intelligible speech. A child with a diagnosis of childhood apraxia of speech specifically experiences problems in programming and planning his/her speech movements.



Speech Plans

When we try to communicate, an idea forms in our brain as to what we want to say. Our brain thinks of words for our messages and puts them in the correct order using correct grammar. This information then translates into a series of highly coordinated movements of the articulators—the lips, jaw, tongue, and soft palate. The brain tells the muscles that control the articulators the

exact order and timing of the movements to produce intelligible speech. Children with apraxia of speech may not be able to accurately and consistently produce speech motor plans. This may be due to neurological impairments or developmental disabilities.



Speech Therapy

When a child receives a diagnosis of CAS, he/she needs to begin speech therapy immediately. Therapy will vary with each child according to the severity of the diagnosis. Experienced SLPs use many different therapy methods since there is not a "one-size-fits-all" program for apraxia. Children with apraxia need frequent one-on-one therapy and lots of repetitive practice with sounds, sequences, and movement patterns in order to incorporate them into their speech and make them automatic. Practicing at home is important to any speech and language program.



Handy Handouts®

Free informational handouts for educators, parents, and students
(cont.)



Practicing Speech

Many therapists recommend the use of sign language, picture communication systems, and other augmentative or alternative communications. This combination of therapies (sign and vocal word attempt) enhances the chance that the listener will be able to understand the speaker. Very young children realize their speech difficulties. Any successful communication experiences will encourage them to work harder. For some children, visual

cues (e.g., pointing to lips for “b” sound) aid in learning to use their articulators to produce a desired word. When the word and sign are consistent, children begin to associate the visual image of the sign with the placement of the articulators.

If you notice speech delays/difficulties with your child’s production or problems with his/her communication skills, please contact your physician or consult a speech-language pathologist in your area.

Related Handy Handouts®:

[50 – Developmental Apraxia of Speech and Developmental Verbal Dyspraxia](#)

[179 – Oral-Motor Workout](#)

[685 – Treatment Methods for Childhood Apraxia of Speech](#)

Resources

Apraxia-KIDS The Childhood Apraxia of Speech Association of North America. Retrieved March 31, 2008, from <http://www.apraxia-kids.org/>

National Institute on Deafness and Other Communication Disorders. Retrieved March 31, 2008, from <http://www.nidcd.nih.gov/health/voice/apraxia.htm>

American Speech-Language Hearing Association (ASHA). Retrieved March 31, 2008, from <http://www.asha.org/public/speech/disorders/ChildhoodApraxia.htm>

Becoming Verbal with Childhood Apraxia. Retrieved March 31, 2008, from www.superduperinc.com

For more FREE Handy Handouts®, go to www.handyhandouts.com